Interferon-gamma immunospot assay of pleural effusion mononuclear cells for diagnosis of tuberculous pleurisy- American Society for Microbiology

DNA probe based colorimetric method for detection of rifampicin resistance of Mycobacterium tuberculosis.- Journal Microbiology Methods

Pyrazinamide susceptibility testing of Mycobacterium tuberculosis by high resolution melt analysis.- Tuberculosis

Rapid Detection of Mycobacterium tuberculosis and Pyrazinamide Susceptibility Related to pncA Mutations in Sputum Specimens through an Integrated Gene-to-Protein Function Approach.-Journal Clinical Microbiology

Evaluation of GeneXpert MTB/RIF for Diagnosis of Tuberculous Meningitis.- Journal Clinical Microbiology

Significant Reduction in Rate of Indeterminate Results of the QuantiFERON-TB Gold In-Tube Test by Shortening Incubation Delay- Journal Clinical Microbiology

Menon Biosensors demonstrates industry-leading virus detection

Standard Operating Procedure (SOP) Specimen processing of CSF, lymph nodes and other tissues for Xpert MTB/RIF

Do We Need to Detect Isoniazid Resistance in Addition to Rifampicin Resistance in Diagnostic Tests for Tuberculosis?- PLoSE ONE

Role of pncA and rpsA Gene Sequencing in Detection of Pyrazinamide Resistance in Mycobacterium tuberculosis Isolates from Southern China. - Journal Clinical Microbiology

Nitrate Reductase Assay for Rapid Detection of Isoniazid, Rifampin, Ethambutol, and Streptomycin Resistance in Mycobacterium tuberculosis: a Systematic Review and Meta-Analysis- Journal Clinical Microbiology

Point-of-care urine test for assessing adherence to isoniazid treatment for tuberculosis.- ERJ

Evaluation of two line probe assays for rapid detection of Mycobacterium tuberculosis, tuberculosis drug resistance, and non-tuberculosis mycobacteria in HIV-infected TB suspects.- Journal Clinical Microbiology

BCG vaccine 'more effective' than thought- Clinical Infectious Disease

Fluoroquinolone resistance in Mycobacterium tuberculosis. What have we learnt? - Int J Tuberc Lung Dis

Tuberculosis drug discovery in the post-postgenomic era- EMBO Molecular Medicine

Improving the health of the tuberculosis drug pipeline- The Lancet

Spectinamides: a new class of semisynthetic antituberculosis agents that overcome native drug

A Trial of Mass Isoniazid Preventive Therapy for Tuberculosis Control- NEJM

Role of clofazimine in the treatment of multidrugresistant tuberculosis: a retrospective observational cohort assessment. - J Antimicrob Chemother.

Evolution and transmission of drug-resistant tuberculosis in a Russian population- Nature

> Sterilization of granulomas is common in active and latent tuberculosis despite within-host variability in bacterial killing- Nature

Mycobacteria manipulate macrophage recruitment through coordinated use of membrane lipids -

The effect of HIV-related immunosuppression on the risk of tuberculosis transmission to household contacts- Clinical Infectious Disease

iNKT Cell Production of GM-CSF Controls Mycobacterium tuberculosis- PLoSE Pathogens

DIAGNOSTIC

VACCINE

DRUGS




Research

Implementation of control

January 2014





Review of the National Tuberculosis Programme in

FIRST JOINT MEETING OF THREE EUROPEAN

Collaborative action on tuberculosis and alcohol

the Republic of Moldova

abuse in Estonia

Global strategy and targets for tuberculosis

Standard Operating Procedure (SOP) Specimen

processing of CSF, lymph nodes and other tissues

Bone marrow stem cells could defeat drug-

When TB Research meets innovative ideas

prevention, care and control after 2015

resistant tuberculosis

ECDC Guidance: investigation and control of TB

incidents affecting children in congregate settings

TUBERCULOSIS NETWORKS

WHO EUROPE

for Xpert MTB/RIF

WHO

TB INNOVATION Blog

ERS/WHO Tuberculosis Consilium: reporting of the

Do high rates of empirical treatment undermine the

Join an online discussion on the Gene Xpert TB

ECDC

TB CARE I - Year 3 Annual Report

tuberculosis in high-burden settings?

diagnostic

potential eff ect of new diagnostic tests for

initial 10 cases.